

SU-ITP-94-14

hep-th/9405103

May 1994

Comment On A Proposal By Strominger[★]LEONARD SUSSKIND[†]*Department of Physics, Stanford University, Stanford, CA 94305-4060*

ABSTRACT

Strominger has proposed an interesting concrete realization of Hawking's idea that information is lost in black hole evaporation. In this note we demonstrate that a straightforward interpretation of Strominger's model leads to a complete breakdown of the conditions for using statistics for analyzing the results of experiments. The probabilities produced by the theory are operationally meaningless.

★ Work supported in part by NSF grant PHY89-17438

† email: susskind@dormouse.stanford.edu

Consider the proposition that the probability for an experiment denoted A to lead to a given result B is $P(A; B)$. This is a meaningless assertion unless the following two conditions are met.

1) The experiment can be performed many times. In particular the initial conditions of the experiment are reproducible. In order to keep the experiments independent they should ideally be performed in widely separated regions of space time.

2) The results of the individual experiments must be statistically independent. This implies that the probability for the outcomes of an ensemble of N identically prepared events to be $B_1, B_2, B_3, \dots B_N$ must be

$$P(A; B_1 \dots B_N) = P(A; B_1)P(A; B_2) \dots P(A; B_N) \quad (1)$$

This requirement can be considered to be a consistency condition on a theory in the following sense. If we regard the ensemble of experiments to itself be a single experiment, the theory can be directly applied to it. Using the same rules of calculation that were used for the single event A , we should directly confirm eq. (1). In ordinary quantum field theory these requirements are insured by the cluster decomposition property.

In recent years the validity of the usual operational rules of quantum mechanics has been questioned principally in the context of black hole formation and evaporation. According to Hawking [1] the process in question inevitably leads to a loss of quantum coherence so that an initial pure state of matter will evolve to a mixed density matrix. It is of obvious interest to ask whether specific versions of Hawking's proposal satisfy the consistency condition of eq. (1). Strominger [2] has put forward one such proposal which is specific enough to analyze from this viewpoint. we shall see that a straightforward interpretation of this proposal fails the test.

Strominger does not consider 3+1 dimensional gravity. His construction is based on the more easily analyzable 1+1 dimensional models of Callen Giddings Harvey and Strominger. It is argued that one particular version of the CGHS model is isomorphic to an open string theory. The physical space- time is identified with the world sheet of a semi-infinite string. The target space of string theory is the space of fields that propagate on the world sheet. According to the model, black hole formation and evaporation is always accompanied by the splitting off of a piece of space which is represented by an open string. The emitted open string carries off the information that fell into the black hole.

Before considering the implications of Strominger's model one thing needs to be clarified. In this paper the terms, experiments and their outcome, refer to operations carried out in the usual way by observers living in the universe. We are not interested in the rules for "superobservers" who perform experiments involving ensembles of universes. We are interested in a set of rules to be used by future experimental physicists like ourselves.

Consider an event in which matter in an initial state A creates a black hole which subsequently evaporates producing an open string in state i and final state B of the semi-infinite universe. The amplitude for this transition is

$$S(A|B, i) \tag{2}$$

The final state of the semi-infinite universe will not generally be pure. According to the most straightforward rule it will be described by the density matrix

$$\sum_i S^*(A|B, i) S(A|B', i) \tag{3}$$

Eq.[(3)] can be pictorially represented as in Fig (1) in which the two sides represent S and S^*




FIGURE 1. The density matrix. The white region represents the world sheet. The shaded region is not part of the universe.

An experiment to determine if information is lost can not be carried out with a single event. We are therefore led to consider an event in which the initial conditions A are replicated many times at widely separated times. For simplicity consider just two copies of the incident state. The event is illustrated in Fig(2).




FIGURE 2. The density matrix for two identical events.

The two emitted open strings are emitted at widely separated world-sheet times but in identical target space configurations. In string theory they would be considered as identical particles and subject to the rules of either Fermi-Dirac or Bose-Einstein statistics. The implication is that in squaring the amplitude and summing over the open strings, two distinct ways of making the identification must be made. This is shown in Fig (3a) and (3b).




FIGURE 3a. First way of identifying open strings.

FIGURE 3b. Second way of identifying open strings; in this figure the cross-hatched region is not part of the worldsheet.

Thus the final density matrix is

$$\begin{aligned}
 & S(A|B, i) S^*(A|B', i) S(A|C, j) S^*(A|C', j) \\
 & \pm S(A|B, i) S^*(A|C', i) S(A|C, j) S^*(A|B', j)
 \end{aligned} \tag{4}$$

(The plus and minus refer to the target space statistics)

The first term in eq. [4] is all that would be expected if the events were

statistically independent. As an example of the peculiar features of eq. [(4)] suppose in both experiments we post select the same final state B. In the theory of a single event, the probability is

$$\sum_i S(A|B, i) S^*(A|B, i) = P_A(B) \quad (5)$$

For the double event the probability obtained from eq. [(4)] is

$$P_A(B, B) = P_A(B)^2 \pm P_A(B)^2 \quad (6)$$

For example if the open strings are target space fermions the double event can not occur at all while for bosons the probability is double the value required by eq. [(5)] . The point is not that we have encountered some unusual behavior of probabilities but that the very conditions for the use of statistics have broken down. We emphasize again that unless experiments can be performed repeatedly, under identical circumstances with statistically independent results there is no meaning to the probability of a given outcome.

It is interesting to see what happens if we try to repair the damage by making a rule which would only allow identification of strings if they are emitted at similar world sheet times. In other words suppose the open strings carry some memory of when they are emitted and can only be treated as identical if emitted at nearby times. In this case there would be a tight correlation between the times that the branching processes take place on the two sides of the figures representing the evolution of the density matrix. This leads to exactly the situation discussed by Banks, Peskin and Susskind [3] and therefore to non conservation of energy.

It is possible that the rules of string theory may have a less straightforward interpretation, perhaps along the lines of Coleman's wormhole calculus. How such a reinterpretation would translate into a set of rules useful to ordinary observers is not clear.

Acknowledgements:

The author thanks Andy Strominger for a preliminary copy of Ref [2]

REFERENCES

1. S. W. Hawking, *Phys. Rev.* **D14** (1976), 2460.
2. A. Strominger, *Black Holes and Open Strings*, preprint UCSBTH-94-13, May 1994, hep-th/9405094.
3. T. Banks, L. Susskind , and M. E. Peskin, *Nucl. Phys.* **B244** (1984), 125.